

liteal catheters for postoperative analgesia after hallux valgus repair [J]. *Anesth Analg*, 2006, 102:258-262.

[11] Morin AM, Eberhart LH, Behnke HK, et al. Does femoral nerve catheter placement with stimulating catheters improve effective placement? A randomized, controlled, and observer-blinded trial [J]. *Anesth Analg*, 2005, 100:1503-1510.

[12] Hayek SM, Ritchey RM, Sessler D, et al. Continuous femoral nerve analgesia after unilateral total knee arthroplasty: stimulating *versus* nonstimulating catheters [J]. *Anesth Analg*, 2006, 103:1565-1570.

[13] Barrington MJ, Olive DJ, McCutcheon CA, et al. Stimulating catheters for continuous femoral nerve blockade after total knee arthroplasty: a randomized, controlled, double-blinded trial [J]. *Anesth Analg*, 2008, 106:1316-1321.

[14] Dauri M, Sidiropoulou T, Fabbi E, et al. Efficacy of continuous femoral nerve block with stimulating catheters *versus* nonstimulating catheters for anterior cruciate ligament reconstruction [J]. *Reg Anesth Pain Med*, 2007, 32:282-287.

[15] Pham DC, Lelong A, Guilley J. Effect on neurostimulation of injectates used for perineural space expansion before placement of a stimulating catheter: normal saline *versus* dextrose 5% in water [J]. *Reg Anesth Pain Med*, 2009, 34:398-403.

[16] Aveline C, Le Roux A, Le Hetet H, et al. Postoperative efficacies of femoral nerve catheters sited using ultrasound combined with neurostimulation compared with neurostimulation alone for total knee arthroplasty [J]. *Eur J Anaesthesiol*, 2010, 27:978-984.

[17] Gucev G, Yasui GM, Chang TY, et al. Bilateral ultrasound-guided continuous ilioinguinal-iliohypogastric block for pain relief after cesarean delivery [J]. *Anesth Analg*, 2008, 106:1220-1222.

[18] Koscielniak-Nielsen ZJ, Rasmussen H, Hesselbjerg L. Long-axis ultrasound imaging of the nerves and advancement of perineural catheters under direct vision: a preliminary report of four cases [J]. *Reg Anesth Pain Med*, 2008, 33:477-482.

[19] Wang AZ, Gu LL, Zhou QH, et al. Ultrasound-guided continuous femoral nerve block for analgesia after total knee arthroplasty: catheter perpendicular to the nerve *versus* catheter parallel to the nerve [J]. *Reg Anesth Pain Med*, 2010, 35:127-131.

[20] Gerhard B, Hartmut B, Hugo VA. Postoperative analgesia after knee surgery: a comparison of three different concentrations of ropivacaine for continuous femoral nerve blockade [J]. *Anesth Analg*, 2007, 105:256-262.

[21] Tang S, Hong X, Huang YG, et al. Sufentanil does not enhance the efficacy of ropivacaine in combined lumbar plexus and sciatic block: A controlled, randomized clinical trial [J]. *Acute Pain*, 2007, 9:59-63.

(收稿日期: 2011-03-16)

• 医学新闻 •

北京协和医院朱兰教授、张烜教授荣获
2011—2012 年度卫生部有突出贡献中青年专家称号

根据国家卫生和计划生育委员会文件（卫人发[2012] 84 号）通知，北京协和医院妇产科朱兰教授、风湿免疫科张烜教授荣获 2011—2012 年度卫生部有突出贡献中青年专家称号。

“卫生部有突出贡献中青年专家” 称号由单位推荐、各省级卫生行政部门初选和国家卫生和计划

生育委员会评审产生，获得此项称号的同志是医药卫生工作者的优秀代表，为推动我国卫生事业发展作出过突出贡献。

（北京协和医院人力资源处）