

- macular oedema (IBEME study) [J]. Br J Ophthalmol, 2008, 92:76-80.
- [23] Elman MJ, Aiello LP, Beck RW, et al. Randomized trial evaluating ranibizumab plus prompt or deferred laser or triamcinolone plus prompt laser for diabetic macular edema [J]. Ophthalmology, 2010, 117:1064-1077.
- [24] Nguyen QD, Shah SM, Heier JS, et al. Primary End Point (Six Months) Results of the Ranibizumab for Edema of the macula in diabetes (READ-2) study [J]. Ophthalmology, 2009, 116:2175-2181.
- [25] Elman MJ, Bressler NM, Qin H, et al. Expanded 2-year follow-up of ranibizumab plus prompt or deferred laser or triamcinolone plus prompt laser for diabetic macular edema [J]. Ophthalmology, 2011, 118:609-614.
- [26] Mirshahi A, Roohipour R, Lashay A, et al. Bevacizumab-augmented retinal laser photocoagulation in proliferative diabetic retinopathy: a randomized double-masked clinical trial [J]. Eur J Ophthalmol, 2008, 18:263-269.
- [27] Filho JA, Messias A, Almeida FP, et al. Panretinal photocoagulation (PRP) versus PRP plus intravitreal ranibizumab for high-risk proliferative diabetic retinopathy [J]. Acta Ophthalmol, 2011, 89:e567-e572.
- [28] Cho WB, Oh SB, Moon JW, et al. Panretinal photocoagulation combined with intravitreal bevacizumab in high-risk proliferative diabetic retinopathy [J]. Retina, 2009, 29:516-522.
- [29] Jorge R, Costa RA, Calucci D, et al. Intravitreal bevacizumab (Avastin) for persistent new vessels in diabetic retinopathy (IBEPE study) [J]. Retina, 2006, 26:1006-1013.
- [30] Erdol H, Turk A, Akyol NN, et al. The results of intravitreal bevacizumab injections for persistent neovascularizations in proliferative diabetic retinopathy after photocoagulation therapy [J]. Retina, 2010, 30:570-577.
- [31] Tonello M, Costa RA, Almeida FP, et al. Panretinal photocoagulation versus PRP plus intravitreal bevacizumab for high-risk proliferative diabetic retinopathy (IBeHi study) [J]. Acta Ophthalmol, 2008, 86:385-389.
- [32] Huang YH, Yeh PT, Chen MS, et al. Intravitreal bevacizumab and panretinal photocoagulation for proliferative diabetic retinopathy associated with vitreous hemorrhage [J]. Retina, 2009, 29:1134-1140.
- [33] Moradian S, Ahmadi H, Malihi M, et al. Intravitreal bevacizumab in active progressive proliferative diabetic retinopathy [J]. Graefes Arch Clin Exp Ophthalmol, 2008, 246:1699-1705.

(收稿日期: 2013-02-28)

· 医学新闻 ·

协和妇产科沈铿教授当选第四届“首都十大健康卫士”

由北京市卫生局、首都卫生系统精神文明建设协调委员会组织开展的第四届“首都十大健康卫士”评选揭晓,我国著名妇科肿瘤专家,北京协和医院妇产科常务副主任沈铿教授当选。

参加本届“首都十大健康卫士”评选的共有57名优秀医务工作者,分别来自中央、部队、武警、高校、厂矿以及北京市、区卫生局所属的驻京医疗卫生机构。经初评评选出前30名候选人,在《北京日报》上刊登照片和事迹,同时,在搜狐网

健康频道、好大夫在线、北京卫生信息网进行投票并公示。期间,受到社会各界的广泛关注,广大网民积极响应,踊跃投票。2013年2月28日,首都卫生系统精神文明协调委员会召开专题会议,审议通过了第四届“首都十大健康卫士”名单,包括沈铿教授在内的10位专家当选。

(北京协和医院党委综合办)