

- insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor- α , overexpressed in human fat cells from insulin-resistant subjects [J]. *J Biol Chem*, 2003, 278: 45777-45784.
- [22] Senn JJ, Klover PJ, Nowak IA, et al. Interleukin-6 induces cellular insulin resistance in hepatocytes [J]. *Diabetes*, 2002, 51: 3391-3399.
- [23] Senn JJ, Klover PJ, Nowak IA, et al. Suppressor of cytokine signaling-3 (SOCS-3), a potential mediator of interleukin-6-dependent insulin resistance in hepatocytes [J]. *J Biol Chem*, 2003, 278: 13740-13746.
- [24] Bastard JP, Jardel C, Bruckert E, et al. Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss [J]. *J Clin Endocrinol Metab*, 2000, 85: 3338-3342.
- [25] Wang X, Bao W, Liu J, et al. Inflammatory markers and risk of type 2 diabetes: a systematic review and meta-analysis [J]. *Diabetes Care*, 2013, 36: 166-175.
- [26] Hotamisligil GS, Budavari A, Murray D, et al. Reduced tyrosine kinase activity of the insulin receptor in obesity-diabetes. Central role of tumor necrosis factor- α [J]. *J Clin Invest*, 1994, 94: 1543-1549.
- [27] Kern PA, Saghizadeh M, Ong JM, et al. The expression of tumor necrosis factor in human adipose tissue. Regulation by obesity, weight loss, and relationship to lipoprotein lipase [J]. *J Clin Invest*, 1995, 95: 2111-2119.
- [28] Ziccardi P, Nappo F, Giugliano G, et al. Reduction of inflammatory cytokine concentrations and improvement of endothelial functions in obese women after weight loss over one year [J]. *Circulation*, 2002, 105: 804-809.
- [29] Hivert MF, Sullivan LM, Fox CS, et al. Associations of adiponectin, resistin, and tumor necrosis factor- α with insulin resistance [J]. *J Clin Endocrinol Metab*, 2008, 93: 3165-3172.
- [30] Lumeng CN, Bodzin JL, Saltiel AR. Obesity induces a phenotypic switch in adipose tissue macrophage polarization [J]. *J Clin Invest*, 2007, 117: 175-184.
- [31] Lee HM, Kim JJ, Kim HJ, et al. Upregulated NLRP3 inflammasome activation in patients with type 2 diabetes [J]. *Diabetes*, 2013, 62: 194-204.
- [32] Zheng Y, Zhang D, Zhang L, et al. Variants of NLRP3 gene are associated with insulin resistance in Chinese Han population with type-2 diabetes [J]. *Gene*, 2013, 530: 151-154.
- [33] Dehghan A, Dupuis J, Barbalic M, et al. Meta-analysis of genome-wide association studies in > 80 000 subjects identifies multiple loci for C-reactive protein levels [J]. *Circulation*, 2011, 123: 731-738.
- [34] Lamkanfi M, Mueller JL, Vitari AC, et al. Glyburide inhibits the Cryopyrin/Nalp3 inflammasome [J]. *J Cell Biol*, 2009, 187: 61-70.

(收稿日期: 2015-07-19)

· 读者 · 作者 · 编者 ·

本刊对来稿中参考文献著录的有关要求

本刊目前执行 GB/T 7714-2005 《文后参考文献著录规则》中有关参考文献著录的规定。参考文献采用顺序编码制著录, 按照引文先后顺序, 用阿拉伯数字连续编号, 在正文引出处加方括号标注于句尾右上角, 或标注于引文作者的右上角。参考文献引用仅限作者亲自阅读过的主要文献, 作者须仔细核对原文以保证文字和原意无误, 同时保证文献著录的各项内容准确。

本刊参考文献著录格式如下:

[期刊] 作者 (3 名及 3 名以内全部著录; 超过 3 名仅著录前 3 名, 后加 “, 等” 或 “, et al”). 题名 [文献类型标志]. 刊名, 年, 卷: 起页-止页.

[电子期刊] 作者 (3 名及 3 名以内全部著录; 超过 3 名仅著录前 3 名, 后加 “, 等” 或 “, et al”). 题名 [文献类型标志/文献载体标志]. 刊名, 年, 卷: 起页-止页 [引用日期]. 获取和访问路径.

[专著] 作者. 书名 [文献类型标志]. 其他责任者 (如翻译者). 版本 (第 1 版不著录). 出版地: 出版者, 出版年: 引文起页-止页.

[专著中析出文献] 析出文献作者. 题名 [文献类型标志] // 专著主要作 (编) 者. 专著名. 版本 (第 1 版不著录). 出版地: 出版者, 出版年: 析出文献起页-止页.